


What we claim is:

1. A composition of matter having the formula:



wherein $R^1 = \text{CH}_2\text{OH}$, $-\text{CONR}_5\text{R}_6$;

- 5 R^3 is selected from the group consisting of C_{1-15} alkyl, halo, NO_2 , CF_3 , CN , OR^{20} , SR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{SO}_2\text{NR}^{20}\text{COR}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{N}(\text{R}^{20})_2\text{NR}^{20}\text{COR}^{22}$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{C}(\text{NR}^{20})\text{NHR}^{23}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{CONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{OCONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{OC}(\text{O})\text{R}^{20}$, $\text{C}(\text{O})\text{OCH}_2\text{OC}(\text{O})\text{R}^{20}$, and $\text{OCON}(\text{R}^{20})_2$, $-\text{CONR}^7\text{R}^8$, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, aryl, heterocyclyl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN , OR^{20} , SR^{20} , $\text{N}(\text{R}^{20})_2$, $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, $\text{SO}_2\text{NR}^{20}\text{COR}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{N}(\text{R}^{20})_2\text{NR}^{20}\text{COR}^{22}$, $\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{C}(\text{NR}^{20})\text{NHR}^{23}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{CONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, $\text{SO}_2\text{NR}^{20}\text{CO}_2\text{R}^{22}$, $\text{OCONR}^{20}\text{SO}_2\text{R}^{22}$, $\text{OC}(\text{O})\text{R}^{20}$, $\text{C}(\text{O})\text{OCH}_2\text{OC}(\text{O})\text{R}^{20}$, and $\text{OCON}(\text{R}^{20})_2$ and wherein optional heteroaryl, aryl, and heterocyclyl substituent is optionally substituted with halo, NO_2 , alkyl, CF_3 , amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, NCOR^{22} , $\text{NR}^{20}\text{SO}_2\text{R}^{22}$, COR^{20} , CO_2R^{20} , $\text{CON}(\text{R}^{20})_2$, $\text{NR}^{20}\text{CON}(\text{R}^{20})_2$, $\text{OC}(\text{O})\text{R}^{20}$, $\text{OC}(\text{O})\text{N}(\text{R}^{20})_2$, SR^{20} , $\text{S}(\text{O})\text{R}^{22}$, SO_2R^{22} , $\text{SO}_2\text{N}(\text{R}^{20})_2$, CN , and OR^{20} ;

R^5 and R^6 are each individually selected from H, C_1 - C_{15} alkyl optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$, $SO_2NR^{20}CO_2R^{22}$, $SO_2NR^{20}CON(R^{20})_2$, $N(R^{20})_2NR^{20}COR^{22}$, $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, $NR^{20}C(NR^{20})NHR^{23}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $CONR^{20}SO_2R^{22}$, $NR^{20}SO_2R^{22}$, $SO_2NR^{20}CO_2R^{22}$, $OCONR^{20}SO_2R^{22}$, $OC(O)R^{20}$, $C(O)OCH_2OC(O)R^{20}$, and $OCON(R^{20})_2$ and wherein optional heteroaryl, aryl, and heterocyclyl substituent is optionally substituted with halo, NO_2 , alkyl, CF_3 , amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, $NCOR^{22}$, $NR^{20}SO_2R^{22}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $NR^{20}CON(R^{20})_2$, $OC(O)R^{20}$, $OC(O)N(R^{20})_2$, SR^{20} , $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, CN, and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl and heteroaryl, wherein the alkyl, alkenyl, alkynyl, aryl, heterocyclyl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, NO_2 , heterocyclyl, aryl, heteroaryl, CF_3 , CN, OR^{20} , SR^{20} , $N(R^{20})_2$, $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, $SO_2NR^{20}COR^{22}$, $SO_2NR^{20}CO_2R^{22}$, $SO_2NR^{20}CON(R^{20})_2$, $N(R^{20})_2NR^{20}COR^{22}$, $NR^{20}CO_2R^{22}$, $NR^{20}CON(R^{20})_2$, $NR^{20}C(NR^{20})NHR^{23}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $CONR^{20}SO_2R^{22}$, $NR^{20}SO_2R^{22}$, $SO_2NR^{20}CO_2R^{22}$, $OCONR^{20}SO_2R^{22}$, $OC(O)R^{20}$, $C(O)OCH_2OC(O)R^{20}$ and $OCON(R^{20})_2$ and wherein optional heteroaryl, aryl and heterocyclyl substituent is optionally substituted with halo, NO_2 , alkyl, CF_3 , amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, $NCOR^{22}$, $NR^{20}SO_2R^{22}$, COR^{20} , CO_2R^{20} , $CON(R^{20})_2$, $NR^{20}CON(R^{20})_2$, $OC(O)R^{20}$, $OC(O)N(R^{20})_2$, SR^{20} , $S(O)R^{22}$, SO_2R^{22} , $SO_2N(R^{20})_2$, CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen, C_{1-15} alkyl, C_{2-15} alkenyl, C_{2-15} alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, aryl, heterocyclyl, and heteroaryl substituents are optionally substituted with from 1 to 3

substituents independently selected from the group consisting of halo, NO₂, heterocyclyl, aryl, heteroaryl, CF₃, CN, OR²⁰, SR²⁰, N(R²⁰)₂, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, SO₂NR²⁰COR²², SO₂NR²⁰CO₂R²², SO₂NR²⁰CON(R²⁰)₂, N(R²⁰)₂ NR²⁰COR²², NR²⁰CO₂R²², NR²⁰CON(R²⁰)₂, NR²⁰C(NR²⁰)NHR²³, COR²⁰, CO₂R²⁰, CON(R²⁰)₂, CONR²⁰SO₂R²², NR²⁰SO₂R²², SO₂NR²⁰CO₂R²², OCONR²⁰SO₂R²², OC(O)R²⁰, C(O)OCH₂OC(O)R²⁰, and OCON(R²⁰)₂ and wherein each optional heteroaryl, aryl, and heterocyclyl substituent is optionally substituted with halo, NO₂, alkyl, CF₃, amino, mono- or di- alkylamino, alkyl or aryl or heteroaryl amide, NCOR²², NR²⁰SO₂R²², COR²⁰, CO₂R²⁰, CON(R²⁰)₂, NR²⁰CON(R²⁰)₂, OC(O)R²⁰, OC(O)N(R²⁰)₂, SR²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, CN, and OR²⁰;

R²⁰ is selected from the group consisting of H, C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from halo, alkyl, mono- or dialkylamino, alkyl or aryl or heteroaryl amide, CN, O-C₁₋₆ alkyl, CF₃, aryl, and heteroaryl;

R²² is selected from the group consisting of C₁₋₁₅ alkyl, C₂₋₁₅ alkenyl, C₂₋₁₅ alkynyl, heterocyclyl, aryl, and heteroaryl, wherein the alkyl, alkenyl, alkynyl, heterocyclyl, aryl, and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from halo, alkyl, mono- or dialkylamino, alkyl or aryl or heteroaryl amide, CN, O-C₁₋₆ alkyl, CF₃, aryl, and heteroaryl; and

wherein R² and R⁴ are selected from the group consisting of H, C₁₋₆ alkyl and aryl optionally substituted with halo, CN, CF₃, OR²⁰ and N(R²⁰)₂, with the proviso that when R² is not hydrogen then R⁴ is hydrogen, and when R⁴ is not hydrogen then R² is hydrogen.

2. The composition of claim 1 wherein R³ is selected from the group consisting of C₁₋₁₅ alkyl, halo, CF₃, CN, OR²⁰, SR²⁰, S(O)R²², SO₂R²², SO₂N(R²⁰)₂, COR²⁰, CO₂R²⁰, -CONR⁷R⁸, aryl and heteroaryl wherein the alkyl, aryl and heteroaryl substituents are

optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF_3 , CN, OR^{20} , SR^{20} , S(O)R^{22} , SO_2R^{22} , $\text{SO}_2\text{N(R}^{20})_2$, COR^{20} , CO_2R^{20} and $\text{CON(R}^{20})_2$, and wherein each optional heteroaryl and aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, and OR^{20} ;

5 R^5 and R^6 are each individually selected from the group consisting of H, and $\text{C}_1\text{-C}_{15}$ alkyl optionally substituted with one aryl substituent that is optionally substituted with halo or CF_3 ;

R^7 is selected from the group consisting of C_{1-15} alkyl, C_{2-15} alkynyl, aryl, and heteroaryl, wherein the alkyl, alkynyl, aryl, and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF_3 , CN, and OR^{20} , and wherein each optional heteroaryl and aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen and C_{1-15} alkyl;

R^{20} is selected from the group consisting of H, C_{1-4} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with one alkyl substituent; and

R^{22} is selected from the group consisting of C_{1-4} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with from 1 to 3 alkyl groups.

3. The composition of claim 1 wherein R^3 is selected from the group consisting of C_{1-15} alkyl, halo, CF_3 , CN, OR^{20} , CO_2R^{20} , $-\text{CONR}^7\text{R}^8$, aryl and heteroaryl, wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, aryl, CF_3 , CN, OR^{20} , CO_2R^{20} or $\text{CON(R}^{20})_2$, and wherein each optional heteroaryl and aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, and OR^{20} ;

R^5 and R^6 are each individually selected from hydrogen and C_{1-6} alkyl;

R^7 is selected from the group consisting of C_{1-10} alkyl, aryl, and heteroaryl, wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF_3 , CN, and OR^{20} , and wherein each optional heteroaryl and aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from the group consisting of hydrogen and C_{1-15} alkyl; and

R^{20} is selected from the group consisting of hydrogen and C_{1-4} alkyl.

4. The composition of claim 1 wherein R^3 is selected from the group consisting of C_{1-10} , alkyl, halo, CF_3 , CN, CO_2R^{20} , $-CONR^7R^8$, aryl and heteroaryl wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, CF_3 , CN, OR^{20} and $CON(R^{20})_2$;

R^5 and R^6 are each individually selected from hydrogen and C_{1-6} alkyl;

R^7 is selected from the group consisting of C_{1-10} alkyl, aryl, and heteroaryl, wherein the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF_3 , CN, OR^{20} and wherein each optional heteroaryl and aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from hydrogen and C_{1-15} alkyl; and

R^{20} is selected from hydrogen and C_{1-4} alkyl.

5. The composition of claim 1 wherein R^3 is selected from the group consisting of C_{1-10} alkyl, halo, CF_3 , CN, OR^{20} , CO_2R^{20} , $-CONR^7R^8$ and aryl; wherein the alkyl and aryl substituents are optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, alkyl, CF_3 , CN, OR^{20} and $CON(R^{20})_2$;

R^5 and R^6 are each individually selected from hydrogen and C_{1-6} ;

R^7 is selected from the group consisting of C_{1-10} alkyl, aryl, and heteroaryl, where the alkyl, aryl and heteroaryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, heteroaryl, CF_3 , CN, OR^{20} and wherein each optional heteroaryl and aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from hydrogen and C_{1-15} alkyl; and

R^{20} is selected from hydrogen and C_{1-4} alkyl.

6. The composition of claim 1 wherein $R^1 = CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$ and aryl; wherein the aryl substituent is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 , CN, OR^{20} , and $CON(R^{20})_2$;

R^7 is selected from the group consisting of hydrogen, C_{1-10} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, CF_3 , CN, OR^{20} and wherein each optional aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, and OR^{20} ;

R^8 is selected from hydrogen and C_{1-15} alkyl; and

R^{20} is selected from hydrogen and C_{1-4} alkyl.

7. The composition of claim 1 wherein $R^1 = CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$ and aryl wherein the aryl substituent is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, C_{1-8} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with one substituent selected from halo, aryl, CF_3 , CN, and OR^{20} and wherein each optional aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN, or OR^{20} ;

R^8 is selected from hydrogen and C_{1-8} alkyl; and

R^{20} is selected from hydrogen and C_{1-4} alkyl.

8. The composition of claim 1 wherein $R^1 = CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$, and aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group of halo, C_{1-3} alkyl, CF_3 , and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, C_{1-8} alkyl that is optionally substituted with one substituent selected from halo, CF_3 , CN and OR^{20} ;

R^8 is selected from hydrogen and C_{1-3} alkyl; and

R^{20} is selected from hydrogen and C_{1-4} alkyl.

9. The composition of claim 1 wherein $R^1 = CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$, and aryl that is optionally substituted with one substituent selected from the group of halo, C_{1-3} alkyl, and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, and C_{1-3} alkyl that is optionally substituted with one substituent selected from halo, CF_3 , CN and OR^{20} ;

R^8 is hydrogen; and

R^{20} is selected from hydrogen and C_{1-4} alkyl.

10. The composition of claim 1 wherein $R^1 = CH_2OH$;

R^3 is selected from the group consisting of CO_2R^{20} , $-CONR^7R^8$, and aryl that is optionally substituted with one substituent selected from halo, C_{1-3} alkyl and OR^{20} ;

R^7 is selected from of hydrogen, and C_{1-3} alkyl;

R^8 is hydrogen; and

R^{20} is selected from hydrogen and C_{1-4} alkyl.

11. The composition of claim 10 wherein R^7 is a methyl.

12. The composition of claim 10 wherein R_3 is $-\text{CO}_2\text{Et}$.

13. The composition of claim 1 wherein $R^1 = -\text{CONHET}$;

R^3 is selected from the group consisting of CO_2R^{20} , $-\text{CONR}^7R^8$, and aryl; that is optionally substituted with from 1 to 3 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 , CN , OR^{20} , and $\text{CON}(R^{20})_2$;

R^7 is selected from the group consisting of hydrogen, C_{1-10} alkyl and aryl, wherein the alkyl and aryl substituents are optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, aryl, CF_3 , CN , and OR^{20} and wherein each optional aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN , and OR^{20} ;

R^8 is selected from hydrogen, and C_{1-15} alkyl; and

R^{20} is selected from hydrogen, and C_{1-4} alkyl.

14. The composition of claim 1 wherein $R^1 = -\text{CONHET}$;

R^3 is selected from the group consisting of CO_2R^{20} , $-\text{CONR}^7R^8$, aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group consisting of halo, C_{1-6} alkyl, CF_3 and OR^{20} ;

R^7 is selected from the group consisting of hydrogen, C_{1-8} alkyl, and aryl, wherein the alkyl and aryl substituents are optionally substituted with one substituent selected from the group consisting of halo, aryl, CF_3 , CN , OR^{20} and each optional aryl substituent is optionally substituted with halo, alkyl, CF_3 , CN , and OR^{20} ;

R^8 is selected from hydrogen, and C_{1-8} alkyl; and

R^{20} is selected from hydrogen, and C_{1-4} alkyl.

15. The composition of claim 1 wherein $R^1 = -\text{CONHET}$;

R^3 is selected from the group consisting of CO_2R^{20} , $-\text{CONR}^7R^8$, and aryl that is optionally substituted with from 1 to 2 substituents independently selected from the group

consisting of halo, C₁₋₃ alkyl, CF₃ and OR²⁰;

R⁷ is selected from the group consisting of hydrogen, and C₁₋₈ alkyl that is optionally substituted with one substituent selected from halo, CF₃, CN and OR²⁰;

R⁸ is selected from hydrogen, and C₁₋₃ alkyl; and

5 R²⁰ is selected from hydrogen, and C₁₋₄ alkyl.

16. The composition of claim 1 wherein R¹ = -CONHET;

R³ is selected from the group consisting of CO₂R²⁰, -CONR⁷R⁸, and aryl that is optionally substituted with one substituent selected from halo, C₁₋₃ alkyl and OR²⁰;

10 R⁷ is selected from the group consisting of hydrogen and C₁₋₃ alkyl that is optionally substituted with one substituent selected from halo, CF₃, CN and OR²⁰;

R⁸ is hydrogen; and

R²⁰ is selected from hydrogen, and C₁₋₄ alkyl.

17. The composition of claim 1 wherein R¹ = -CONHET;

15 R³ is selected from the group consisting of CO₂R²⁰, -CONR⁷R⁸, and aryl that is optionally substituted with one substituent selected from halo, C₁₋₃ alkyl and OR²⁰;

R⁷ is selected from hydrogen, and C₁₋₃ alkyl;

R⁸ is hydrogen; and

R²⁰ is selected from hydrogen, and C₁₋₄ alkyl.

18. The composition of claim 10 where R¹ is -CONHET.

20 19. A composition matter of claim 1 wherein the compound is selected from
ethyl-1-{9-[(4S,2R,3R,5R)-3,4-dihydroxy-5-(hydroxymethyl)oxolan-2-yl]-6-aminopurin-2-
yl}pyrazole-4-carboxylate, (4S,2R,3R,5R)-2-{6-amino-2-[4-(4-chlorophenyl)-
pyrazolyl]purin-9-yl}-5-(hydroxymethyl)oxolane-3,4-diol, (4S,2R,3R,5R)-2-{6-amino-2-[4-
(4-methoxyphenyl)pyrazolyl]purin-9-yl}-5-(hydroxymethyl)oxolane-3,4-diol, (4S,2R,3R,5R)-
25 2-{6-amino-2-[4-(4-methylphenyl)pyrazolyl]purin-9-yl}-5-(hydroxymethyl)-oxolane-3,4-diol,

(1-{9-[(4S,2R,3R,5R)-3,4-dihydroxy-5-(hydroxymethyl)oxolan-2-yl]-6-aminopurin-2-yl}pyrazol-4-yl)-N-methylcarboxamide, 1-{9-[(4S,2R,3R,5R)-3,4-dihydroxy-5-(hydroxymethyl)oxolan-2-yl]-6-aminopurin-2-yl}pyrazole-4-carboxylic acid, and mixtures thereof.

Sub 4
20. A method for stimulating coronary vasodilatation in a mammal by administering to the mammal a therapeutically effective amount of a compound of claim 1 that is sufficient to stress the heart and induce a coronary steal situation for the purposes of imaging the heart.

12 21. The method of claim 20 wherein the therapeutically effective amount ranges from about 0.01 to about 100 mg/kg weight of the mammal.

10 13 22. The method of claim 20 wherein the mammal is a human.

Sub 5
23. A pharmaceutical composition of matter comprising the composition of claim 1 and one or more pharmaceutical excipients;

24. The pharmaceutical composition of matter of claim 23 wherein the pharmaceutical composition is in the form of a solution.

15 25. The pharmaceutical composition of matter of claim 23 wherein the composition is useful as an anti-inflammatory, in adjunctive therapy with angioplasty, as a platelet aggregation inhibitor, and as an inhibitor of platelet and neutrophil activation.

20

add
a6

31

A